



Fagron Advanced Derma

Compatibility Table

API/DCI	Concentration (%)	Emolivan™	Fitalite™	Nourisil™	Nourivan™ Antiox	Occluvan™	Seraqua™	Versatile™	Versatile™ Anhydrous*	Versatile™ Rich	Espumil™
		Beyond-use Date (days)									
17- α -Estradiol	0.025 to 0.05		90				90		180		90
Aloe vera extract	0.5 to 10.0	90	90 (5%)	90	30 (8%)		90	90	180	90	
Alpha-bisabolol	0.5 to 5.0	90	90		30		90 (1%)	90	180	90	
Ammonium lactate	1.0 to 12.0	90			30 (8.0%)			90 (10.0%)	180	90 (10.0%)	
Anthralin/dithranol	0.05 to 3.0	90			30	90		14	180	14	
Arginine HCl	2.5	30			30			90	180	90	
Ascorbic acid (vit. C)	5.0 to 15.0			14	60		14 (10.0%)	60	180	90	
Azelaic acid	10.0 to 20.0		90					90	180	90	
Benzoyl peroxide	2.5 10.0		30 (5%)					30	180	30	
Benzyl benzoate	25.0	90						90	180		
Betamethasone dipropionate	0.05	90	30		30	90		90	180	90	
Betamethasone valerate	0.025 to 0.1	90	30		30	90		90	180	90	
Caffeine	1.0 to 2.0		90		30		90	90	180	90	90
Chamomile extract	0.5 to 5.0	90	90	90	30	90	90	90	180	90	
Ciclopirox olamine	1.0	90	90		30	90		90	180	90	
Clindamycin HCl	1.0 to 3.0		90	90				30	180	60	
Clobetasol propionate	0.05	90	90		30	90	90	90	180	90	60



API/DCI	Concentration (%)	Emolivan™	Fitalite™	Nourisil™	Nourivan™ Antiox	Occluvan™	Seraqua™	Versatile™	Versatile™ Anhydrous*	Versatile™ Rich	Espumil™
		Beyond-use Date (days)									
Clotrimazole	1.0 to 2.0	90	90					90	180	90	
Coal tar	1.0 to 10.0	90				90		90	180	90	
Coal tar solution	5.0 to 20.0	90				14 (10.0%)		90 (10.0%)	180	90 (10.0%)	
Cyanocobalamin (vit. B12)	0.07	90	30		30	90	90	30	180	30	
Desonide	0.05 to 0.1	90	90		30	90		90	180	90	
Desoximetasone	0.25	90	90		30	90		90	180	90	
Dexpanthenol	0.5 to 5.0	90	90	90	30	90	90	90	180	90	
Diclofenac sodium	3.0	90					90	90	180	90	
Diflucortolone valerate	0.1 to 0.3	90	90		30	90		90	180	90	
Erythromycin	0.5 to 4.0		90		30			90	180	90	
Finasteride	0.1						90		180		90
Fluocinolone acetonide	0.01 to 0.1	90	90		30	90		90	180	90	
Glycerol	0.5 to 20.0	90	90 (5.0%)	90	30 (8.0%)	90	90	90	180	90	
Glycolic acid	5.0 to 15.0		90		30		90	90	180	90	
Hyaluronic acid	0.2 to 2.5			30		90			180		
Hydrocortisone	0.25 to 2.5	90	90		90	90		90	180	90	
Hydrocortisone acetate	0.25 to 2.5	90	90		90	90		90	180	90	
Hydroquinone	2.0 to 4.0				60				180		



API/DCI	Concentration (%)	Emolivan™	Fitalite™	Nourisil™	Nourivan™ Antiox	Occluvan™	Seraqua™	Versatile™	Versatile™ Anhydrous*	Versatile™ Rich	Espumil™
		Beyond-use Date (days)									
Ketoconazole	2.0	90	90				90	90	180	90	90
Kojic acid	1.0 to 4.0		30		30		14		180		
Lactic acid	1.0 to 20.0	90 (10%)	90				90	90	180	90	
Latanoprost	0.005		30				120		180		90
Lidocaine	0.5 to 10.0			90		90 (5.0%)			180		
Lidocaine HCl	0.5 to 10.0	14 (5.0%)			30		90	90	180	90	
Melatonin	0.0033 to 0.1		90				90		180		90
Menthol	1.0 to 3.0	90	90	90	30	90	90	90	180	90	90
Metronidazole	0.5 to 3.0		90				90	90	180	90	
Minoxidil	2.0 to 5.0%		90				90		180		180
Mometasone furoate	0.1	90	30		30	90		90	180	90	
Nicotinamide (vit. B3)	2.0 to 5.0	90	90		30	90	90	90	180	90	
Permethrin	5.0	90	90						180	90	
Prostaquinon™	3.0 to 30.0			90					180		90 (3.0%)
Resorcinol	15.0				90				180		
Retinol (vit. A)	0.4		30		0		60	30	180	30	
Salicylic acid	0.5 to 20.0	90	90 (5.0%)			90		90 (10.0%)	180	90 (10.0%)	
Shea butter	1.0 to 20.0	90	30 (5.0%)		30 (8.0%)	90		90	180		
Sodium hyaluronate	1.0				30		90		180		



API/DCI	Concentration (%)	Emolivan™	Fitalite™	Nourisil™	Nourivan™ Antiox	Occluvan™	Seraqua™	Versatile™	Versatile™ Anhydrous*	Versatile™ Rich	Espumil™
		Beyond-use Date (days)									
Sulfur	5.0 to 10.0	90	30 (5.0%)		30 (8.0%)	90		30	180	30	
Tacrolimus	0.03 to 0.5	90	30		30	90		30	180	30	
Tetracycline HCl	2.0 to 3.0				30			90	180	90	
Tocopherol (vit. E)	0.5 to 20.0	14	90 (5.0%)	90 (5.0%)	30 (8.0%)	90	14	90	180	14	
Tretinoin (vit. A acid)	0.01 to 0.1		30		30		90	30	180	30	90
Triamcinolone acetonide	0.025 to 0.1	90	90		30	90		90	180	90	
Urea	5.0 to 40.0	90 (10.0%)	14				90	30	180	60	
Zinc oxide	1.0 to 40.0	90	90 (5.0%)	90 (25.0%)	30 (8.0%)	90		90 (10.0%)	180	90 (25.0%)	
Zinc sulfate	1.0 to 10.0	90	90 (5.0%)		30 (8.0%)	90		90	180	90	

API = Active Pharmaceutical ingredient; DCI = Dermaceutical ingredient.

* Estimated beyond-use date. In the absence of stability information, the USP <795> can be used as a guideline, which advises a BUD of no more than 180 days under refrigeration (2 - 8 °C) or at controlled room temperature (20 - 25 °C) for non-aqueous dosage forms.

Together
we create the future
of personalizing medicine.



Fagron BV
Fascinatio Boulevard 350
3065 WB Rotterdam
The Netherlands

T +31 88 33 11 288
F +31 88 33 11 210
www.fagron.com

 **Fagron**
personalizing
medicine